

References for Hidden Epidemic: Complications of Insufficient Feeding in Breastfed Newborns

By Christie del Castillo-Hegyi

1. Del Castillo-Hegyi C, Achilles J, Segrave-Daly BJ, Hafken L. Fatal Hypernatremic Dehydration in a Term Exclusively Breastfed Newborn. *Children*. 2022;9(9):1379. doi:[10.3390/children9091379](https://doi.org/10.3390/children9091379).
2. Flaherman VJ, Schaefer EW, Kuzniewicz MW, Li SX, Walsh EM, Paul IM. Early weight loss nomograms for exclusively breastfed newborns. *Pediatrics*. 2015;135(1):e16-e23. doi:[10.1542/peds.2014-1532](https://doi.org/10.1542/peds.2014-1532).
3. Marchini G, Persson B, Berggren V, Hagenäs L. Hunger behaviour contributes to early nutritional homeostasis. *Acta Paediatr*. 1998;87(6):671-675. doi:[10.1080/080352598750014094](https://doi.org/10.1080/080352598750014094).
4. Nommsen-Rivers LA, Heinig MJ, Cohen RJ, Dewey KG. Newborn wet and soiled diaper counts and timing of onset of lactation as indicators of breastfeeding inadequacy. *J Hum Lact*. 2008;24(1):27-33. doi:[10.1177/0890334407311538](https://doi.org/10.1177/0890334407311538).
5. Yamane J, Ishibashi S, Sakaguchi M, Kuroiwa T, Kanemura Y, Nakamura M, Miyoshi H, Sawamoto K, Toyama Y, Mizusawa H, Okano H. Transplantation of human neural stem/progenitor cells overexpressing galectin-1 improves functional recovery from focal brain ischemia in the Mongolian gerbil. *Mol Brain*. 2011;4:35. doi:[10.1186/1756-6606-4-35](https://doi.org/10.1186/1756-6606-4-35).
6. Zhou D, Qian J, Liu CX, Chang H, Sun RP. Repetitive and profound insulin-induced hypoglycemia results in brain damage in newborn rats: an approach to establish an animal model of brain injury induced by neonatal hypoglycemia. *Eur J Pediatr*. 2008;167(10):1169-1174. doi:[10.1007/s00431-007-0653-2](https://doi.org/10.1007/s00431-007-0653-2).
7. Auer RN. Hypoglycemic brain damage. *Metab Brain Dis*. 2004;19(3-4):169-175. doi:[10.1023/b:mebr.0000043967.78763.5b](https://doi.org/10.1023/b:mebr.0000043967.78763.5b).
8. Academy of Breastfeeding Medicine Protocol Committee. ABM clinical protocol #22: guidelines for management of jaundice in the breastfeeding infant equal to or greater than 35 weeks' gestation. *Breastfeed Med*. 2010;5(2):87-93. doi:[10.1089/bfm.2010.9994](https://doi.org/10.1089/bfm.2010.9994).
9. Chang RJ, Chou HC, Chang YH, Chen MH, Chen CY, Hsieh WS, Tsao PN. Weight loss percentage prediction of subsequent neonatal hyperbilirubinemia in exclusively breastfed neonates. *Pediatr Neonatol*. 2012;53(1):41-44. doi:[10.1016/j.pedneo.2011.11.008](https://doi.org/10.1016/j.pedneo.2011.11.008).

10. Yang W-C, Zhao L-L, Li Y-C, et al. Bodyweight loss in predicting neonatal hyperbilirubinemia 72 hours after birth in term newborn infants. *BMC Pediatr*. 2013;13:145. doi:[10.1186/1471-2431-13-145](https://doi.org/10.1186/1471-2431-13-145).
11. Bertini G, Dani C, Tronchin M, Rubaltelli FF. Is breastfeeding really favoring early neonatal jaundice? *Pediatrics*. 2001;107(3):e41. doi:[10.1542/peds.107.3.e41](https://doi.org/10.1542/peds.107.3.e41).
12. Ben XM. Nutritional management of newborn infants: practical guidelines. *World J Gastroenterol*. 2008;14(40):6133-6139. doi:[10.3748/wjg.14.6133](https://doi.org/10.3748/wjg.14.6133).
13. Joint FAO/WHO/UNU Expert Consultation. *Human Energy Requirements*. FAO Food and Nutrition Technical Report Series 1. Rome, Italy: Food and Agriculture Organization of the United Nations; 2001. Accessed July 10, 2021. <https://www.fao.org/4/y5686e/y5686e00.htm>.
14. Maintenance fluid: Maintenance Fluid Calculation for Children. *Merck Manual Professional Version*. Accessed July 10, 2021. <https://www.merckmanuals.com/medical-calculators/MaintenanceFluidChildren.htm>.
15. Gidrewicz DA, Fenton TR. A systematic review and meta-analysis of the nutrient content of preterm and term breast milk. *BMC Pediatr*. 2014;14:216. doi:[10.1186/1471-2431-14-216](https://doi.org/10.1186/1471-2431-14-216).
16. Neville MC, Keller R, Seacat J, Lutes V, Neifert M, Casey C, Allen J, Archer P. Studies in human lactation: milk volumes in lactating women during the onset of lactation and full lactation. *Am J Clin Nutr*. 1988;48(6):1375-1386. doi:[10.1093/ajcn/48.6.1375](https://doi.org/10.1093/ajcn/48.6.1375).
17. Kato I, Horike K, Kawada K, et al. The Trajectory of Expressed Colostrum Volume in the First 48 Hours Postpartum: An Observational Study. *Breastfeed Med*. 2022;17(1):52-58. doi:[10.1089/bfm.2020.0366](https://doi.org/10.1089/bfm.2020.0366).
18. Futatani T, Shimao A, Ina S, et al. Capillary Blood Ketone Levels as an Indicator of Inadequate Breast Milk Intake in the Early Neonatal Period. *J Pediatr*. 2017;191:76-81. doi:[10.1016/j.jpeds.2017.08.080](https://doi.org/10.1016/j.jpeds.2017.08.080).
19. Gartner L. Managing jaundice in the breastfeeding infant. Presented at: California Breastfeeding Summit; January 2013. <https://fedisbest.org/2018/01/starvation-jaundice-brain-injury-breastfed-newborns/>
20. Ogbe Z, Andegiorgish AK, Zeraf AH, Zeng L. Neonatal Hyponatremic Dehydration Associated with Lactation Failure. Khashu M, ed. *Case Reports in Critical Care*. 2020;2020:1-5. doi:[10.1155/2020/8879945](https://doi.org/10.1155/2020/8879945).

21. Dewey KG, Nommsen-Rivers LA, Heinig MJ, Cohen RJ. Risk factors for suboptimal infant breastfeeding behavior, delayed onset of lactation, and excess neonatal weight loss. *Pediatrics*. 2003;112(3 Pt 1):607-619. doi:[10.1542/peds.112.3.607](https://doi.org/10.1542/peds.112.3.607).
22. Nommsen-Rivers LA, Chantry CJ, Pearson JM, Cohen RJ, Dewey KG. Delayed onset of lactogenesis among first-time mothers is related to maternal obesity and factors associated with ineffective breastfeeding. *Am J Clin Nutr*. 2010;92(3):574-584. doi:[10.3945/ajcn.2010.29192](https://doi.org/10.3945/ajcn.2010.29192).
23. Preusting I, Brumley J, Odibo L, Spatz DL, Louis JM. Obesity as a predictor of delayed lactogenesis II. *J Hum Lact*. 2017;33(4):684-691. doi:[10.1177/0890334417727716](https://doi.org/10.1177/0890334417727716).
24. Young PC, Korgenski K, Buchi KF. Early readmission of newborns in a large health care system. *Pediatrics*. 2013;131(5):e1538-e1544. doi:[10.1542/peds.2012-2634](https://doi.org/10.1542/peds.2012-2634).
25. Wilson JL, Wilson BH. Is the "breast is best" mantra an oversimplification? *J Fam Pract*. 2018;67(6):E1-E9. Accessed July 10, 2021. <https://www.mdedge.com/jfponline/article/166932/pediatrics/breast-best-mantra-oversimplification>
26. Flaherman V, Schaefer EW, Kuzniewicz MW, Li SX, Walsh EM, Paul IM. Health Care Utilization in the First Month After Birth and Its Relationship to Newborn Weight Loss and Method of Feeding. *Acad Pediatr*. 2018;18(6):677-684. doi:[10.1016/j.acap.2017.11.005](https://doi.org/10.1016/j.acap.2017.11.005).
27. Escobar GJ. et al. Rehospitalization for neonatal dehydration: a nested case-control study. *Arch Pediatr Adolesc Med*. 2002;156(2):155-161. doi:[10.1001/archpedi.156.2.155](https://doi.org/10.1001/archpedi.156.2.155).
28. Dassios T, Greenough A, Leontiadis S, Hickey A, Kametas NA. Admissions for hypoglycaemia after 35 weeks of gestation: perinatal predictors of cost of stay. *J Matern Fetal Neonatal Med*. 2019;32(3):448-454. doi:[10.1080/14767058.2017.1381905](https://doi.org/10.1080/14767058.2017.1381905).
29. Lee BK, Le Ray I, Yu Sun J, Wikman A, Reilly M, Johansson S. Haemolytic and Non-Haemolytic Neonatal Jaundice Have Different Risk Factor Profiles. *Acta Paediatr*. 2016;105(12):1444-1450. doi:[10.1111/apa.13470](https://doi.org/10.1111/apa.13470).
30. Chang R-J, Chou H-C, Chen M-H, et al. Weight loss percentage prediction of subsequent neonatal hyperbilirubinemia in exclusively breastfed neonates. *Pediatr Neonatol*. 2012;53(1):41-44. doi:[10.1016/j.pedneo.2011.11.008](https://doi.org/10.1016/j.pedneo.2011.11.008).

31. Chen C-F, Hsu M-C, Shen C-H, et al. Influence of breast-feeding on weight loss, jaundice, and waste elimination in neonates. *Pediatr Neonatol*. 2011;52(2):85–92. doi:[10.1016/j.pedneo.2011.02.010](https://doi.org/10.1016/j.pedneo.2011.02.010).
32. Han S, Yu Z, Liu L, et al. A Model for Predicting Significant Hyperbilirubinemia in Neonates From China. *Pediatrics*. 2015;136(4):e896-e905. doi:[10.1542/peds.2014-4058](https://doi.org/10.1542/peds.2014-4058).
33. ChenY-J, Chen W-C, Chen C-M. Risk factors for hyperbilirubinemia in breastfed term neonates. *Eur J Pediatr*. 2012;171:167–171. doi:[10.1007/s00431-011-1512-8](https://doi.org/10.1007/s00431-011-1512-8).
34. Huang M-S, Lin M-C, Chen H-H, Chien K-L, Chen C-H. Risk factor analysis for late-onset neonatal hyperbilirubinemia in Taiwanese infants. *Pediatr Neonatol*. 2009;50(6):261-265. doi:[10.1016/S1875-9572\(09\)60074-7](https://doi.org/10.1016/S1875-9572(09)60074-7).
35. Huang A, Tai B-C, Wong L-Y, Lee J, Yong E-L. Differential risk for early breastfeeding jaundice in a multi-ethnic Asian cohort. *Ann Acad Med Singapore*. 2009;38(3):217-224. PMID: [19347075](https://pubmed.ncbi.nlm.nih.gov/19347075/).
36. Tarcan A, Tiker F, Şalk Vatandaş N, Haberal A, Gürakan B. Weight loss and hyponatremia in breast-fed babies: frequency in neonates with non-hemolytic jaundice. *J Paediatr Child Health*. 2005;41(9-10):484-487. doi:[10.1111/j.1440-1754.2005.00688.x](https://doi.org/10.1111/j.1440-1754.2005.00688.x).
37. Kuzniewicz M, Greene DN, Walsh EM, McCulloch CE, Newman TB. Association Between Laboratory Calibration of a Serum Bilirubin Assay, Neonatal Bilirubin Levels, and Phototherapy Use. *JAMA Pediatr*. 2016;170(6):557-556. doi:[10.1001/jamapediatrics.2015.4944](https://doi.org/10.1001/jamapediatrics.2015.4944).
38. Yu T-C, Nguyen C, Ruiz N, et al. Prevalence and burden of illness of treated hemolytic neonatal hyperbilirubinemia in a privately insured population in the United States. *BMC Pediatr*. 2019;19:53. doi:[10.1186/s12887-019-1414-x](https://doi.org/10.1186/s12887-019-1414-x)
39. Maisels MJ, Gifford K, Antle CE, Leib GR. Jaundice in the healthy newborn infant: a new approach to an old problem. *Pediatrics*. 1988;81(4):505-511. PMID: [3353184](https://pubmed.ncbi.nlm.nih.gov/3353184/).
40. Muri JH, Crawford N, Jellen BC. Reducing avoidable obstetrical and neonatal readmissions (presentation slides). American Hospital Association. <http://www.aha.org/content/11/PerinatalReadmissionscall1.pdf>
41. Samayam P, Ranganathan PK, Kotari UD, Balasundaram R. Study of Asymptomatic Hypoglycemia in Full Term Exclusively Breastfed Neonates in First 48 Hours of Life. *J Clin Diagn Res*. 2015;9(9):SC07-SC10. doi:[10.7860/JCDR/2015/14971.6610](https://doi.org/10.7860/JCDR/2015/14971.6610).

42. Harris DL, Weston PJ, Gamble GD, Harding JE. Glucose Profiles in Healthy Term Infants in the First 5 Days: The Glucose in Well Babies (GLOW) Study. *J Pediatr*. 2020;223:34-41.e4. doi:[10.1016/j.jpeds.2020.02.079](https://doi.org/10.1016/j.jpeds.2020.02.079).
43. Oddie SJ, Craven V, Deakin K, Westman J, Scally A. Severe neonatal hypernatraemia: a population based study. *Arch Dis Child Fetal Neonatal Ed*. 2013;98(5):F384-F387. doi:[10.1136/archdischild-2012-302908](https://doi.org/10.1136/archdischild-2012-302908).
44. Laing IA, Wong CM. Hypernatraemia in the first few days: is the incidence rising? *Arch Dis Child Fetal Neonatal Ed*. 2002;87(3):F158-162.
45. Wang AC, Chen SJ, Yuh YS, Hua YM, Lu TJ, Lee CM. Breastfeeding-associated neonatal hypernatremic dehydration in a medical center: a clinical investigation. *Acta Paediatr Taiwan*. 2007;48(4):186-190.
46. Ferrández-González M, Bosch-Giménez V, López-Lozano J, Moreno-López N, Palazón-Bru A, Cortés-Castell E. Weight loss thresholds to detect early hypernatremia in newborns. *J Pediatr (Rio J)*. 2018 Jul 18. pii:S0021-7557(18)30386-3. doi:[10.1016/j.jped.2018.06.005](https://doi.org/10.1016/j.jped.2018.06.005).
47. Musapasaoglu H, Agildere AM, Teksam M, Tarcan A, Gurakan B. Hypernatraemic dehydration in a neonate: brain MRI findings. *Br J Radiol*. 2008;81(962):e57-e60. doi:[10.1259/bjr/28766369](https://doi.org/10.1259/bjr/28766369).
48. Korkmaz A, Yiğit S, Firat M, Oran O. Cranial MRI in neonatal hypernatraemic dehydration. *Pediatr Radiol*. 2000;30(5):323-325. doi:[10.1007/s002470050749](https://doi.org/10.1007/s002470050749).
49. Meena A, Singh A, Goyal VK, Gupta N, Payal V, Chaturvedi K. Brain Injury Patterns in Neonates With Hypernatremic Dehydration: Single Center Experience. *Indian Pediatr*. 2021;58(10):947-950. PMID: [34636326](https://pubmed.ncbi.nlm.nih.gov/34636326/).
50. Koklu E, Gunes T, Ozturk MA, Kose M, Kurtoglu S, Yuksel F. A review of 116 cases of breastfeeding-associated hypernatremia in rural area of central Turkey. *J Trop Pediatr*. 2007;53(5):347-350. doi:[10.1093/tropej/fmm026](https://doi.org/10.1093/tropej/fmm026).
51. Boskabadi H, Akhondian J, Afarideh M, et al. Long-Term Neurodevelopmental Outcome of Neonates with Hypernatremic Dehydration. *Breastfeed Med*. 2017;12(3):163-168. doi:[10.1089/bfm.2016.0054](https://doi.org/10.1089/bfm.2016.0054).
52. Boskabadi H, Zakerihamidi M, Moradi A. Predictability of prognosis of infantile hypernatremic dehydration: a prospective cohort study. *J Matern Fetal Neonatal Med*. 2022;35(1):66-74. doi:[10.1080/14767058.2020.1712698](https://doi.org/10.1080/14767058.2020.1712698).

53. Lavagno C, Camozzi P, Renzi S, et al. Breastfeeding-Associated Hyponatremia: A Systematic Review of the Literature. *J Hum Lact*. 2016;32(1):67-74. doi:[10.1177/0890334415613079](https://doi.org/10.1177/0890334415613079).
54. Kaiser JR, Bai S, Gibson N, et al. Association Between Transient Newborn Hypoglycemia and Fourth-Grade Achievement Test Proficiency: A Population-Based Study. *JAMA Pediatr*. 2015;169(10):913-921. doi:[10.1001/jamapediatrics.2015.1631](https://doi.org/10.1001/jamapediatrics.2015.1631).
55. Wickström R, Skiöld B, Petersson G, Stephansson O, Altman M. Moderate neonatal hypoglycemia and adverse neurological development at 2-6 years of age. *Eur J Epidemiol*. 2018;33(10):1011-1020. doi:[10.1007/s10654-018-0425-5](https://doi.org/10.1007/s10654-018-0425-5).
56. Seske LM, Merhar SL, Haberman BE. Late-Onset Hypoglycemia in Term Newborns With Poor Breastfeeding. *Hosp Pediatr*. 2015;5(9):501-504. doi:[10.1542/hpeds.2015-0086](https://doi.org/10.1542/hpeds.2015-0086).
57. Marwah A, Gathwala G. Symptomatic hypoglycemia causing brain injury in a term breast-fed newborn following early discharge. *Indian J Pediatr*. 2011;78(12):1549-1551. doi:[10.1007/s12098-011-0465-2](https://doi.org/10.1007/s12098-011-0465-2).
58. Burns CM, Rutherford MA, Boardman JP, Cowan FM. Patterns of cerebral injury and neurodevelopmental outcomes after symptomatic neonatal hypoglycemia. *Pediatrics*. 2008;122(1):65-74. doi:[10.1542/peds.2007-2822](https://doi.org/10.1542/peds.2007-2822).
59. Jangaard KA, Fell DB, Dodds L, Allen AC. Outcomes in a population of healthy term and near-term infants with serum bilirubin levels of ≥ 325 micromol/L (≥ 19 mg/dL) who were born in Nova Scotia, Canada, between 1994 and 2000. *Pediatrics*. 2008;122(1):119-124. doi:[10.1542/peds.2007-0967](https://doi.org/10.1542/peds.2007-0967).
60. Bhutani VK, Johnson-Hamerman L. The clinical syndrome of bilirubin-induced neurologic dysfunction. *Semin Fetal Neonatal Med*. 2015;20(1):6-13. doi:[10.1016/j.siny.2014.12.008](https://doi.org/10.1016/j.siny.2014.12.008).
61. Wei C-C, Chang C-H, Lin C-L, Chang S-N, Li T-C, Kao C-H. Neonatal jaundice and increased risk of attention-deficit hyperactivity disorder: a population-based cohort study. *J Child Psychol Psychiatry*. 2015;56(4):460-467. doi:[10.1111/jcpp.12303](https://doi.org/10.1111/jcpp.12303).
62. Ricardo-Garcell J, Harmony T, Porras-Kattz E, et al. Epileptic encephalopathy in children with risk factors for brain damage. *Epilepsy Res Treat*. 2012;2012(1):747565. doi:[10.1155/2012/747565](https://doi.org/10.1155/2012/747565).
63. Andrade E, Araujo Júnior E, Rolo LC, Da Silva Costa F. Risk factors for cerebral palsy in premature infants identified during the pre and perinatal periods: a case-control study. *Minerva Ginecol*. 2016;68(1):29-36. PMID: [25415488](https://pubmed.ncbi.nlm.nih.gov/25415488/).

64. Maimburg RD, Olsen J, Sun Y. Neonatal hyperbilirubinemia and the risk of febrile seizures and childhood epilepsy. *Epilepsy Res.* 2016;124:67-72. doi:[10.1016/j.eplepsyres.2016.05.004](https://doi.org/10.1016/j.eplepsyres.2016.05.004).
65. Wusthoff CJ, Loe IM. Impact of bilirubin-induced neurologic dysfunction on neurodevelopmental outcomes. *Semin Fetal Neonatal Med.* 2015;20(1):52-57. doi:[10.1016/j.siny.2014.12.003](https://doi.org/10.1016/j.siny.2014.12.003).
66. Sarici D, Gunes T, Yazici C, et al. Investigation on malondialdehyde, S100B, and advanced oxidation protein product levels in significant hyperbilirubinemia and the effect of intensive phototherapy on these parameters. *Pediatr Neonatol.* 2015;56(2):95-100. doi:[10.1016/j.pedneo.2014.06.006](https://doi.org/10.1016/j.pedneo.2014.06.006).
67. Okumus N, Turkyilmaz C, Onal EE, et al. Tau and S100B proteins as biochemical markers of bilirubin-induced neurotoxicity in term neonates. *Pediatr Neurol.* 2008;39(4):245-252. doi:[10.1016/j.pediatrneurol.2008.07.004](https://doi.org/10.1016/j.pediatrneurol.2008.07.004).
68. Bhutani VK, Johnson L. Kernicterus in the 21st century: frequently asked questions. *J Perinatol.* 2009;29(Suppl 1):s20-s24. doi:[10.1038/jp.2008.212](https://doi.org/10.1038/jp.2008.212).
69. Hokkanen L, Launes J, Michelsson K. Adult neurobehavioral outcome of hyperbilirubinemia in full term—a 30 year prospective follow-up study. *PeerJ.* 2014;2:e294. doi:[10.7717/peerj.294](https://doi.org/10.7717/peerj.294).
70. Escobar GJ, Liljestrand P, Hudes ES, et al. Five-year neurodevelopmental outcome of neonatal dehydration. *J Pediatr.* 2007;151(2):127-133.e1. doi:[10.1016/j.jpeds.2007.03.009](https://doi.org/10.1016/j.jpeds.2007.03.009).
71. Thornton PS, Stanley CA, De Leon DD, et al. Recommendations from the Pediatric Endocrine Society for Evaluation and Management of Persistent Hypoglycemia in Neonates, Infants, and Children. *J Pediatr.* 2015;167(2):238-245. doi:[10.1016/j.jpeds.2015.03.057](https://doi.org/10.1016/j.jpeds.2015.03.057).
72. Wickremasinghe AC, Kuzniewicz MW, McCulloch CE, Newman TB. Efficacy of Subthreshold Newborn Phototherapy During the Birth Hospitalization in Preventing Readmission for Phototherapy. *JAMA Pediatr.* 2018;172(4):378-385. doi:[10.1001/jamapediatrics.2017.5630](https://doi.org/10.1001/jamapediatrics.2017.5630).
73. Zaitseva M, Yoshihara T, Nakai H, Kubota S. Optimal Thermal Control with Sufficient Nutrition May Reduce the Incidence of Neonatal Jaundice by Preventing Body-Weight Loss Among Non-Low Birth Weight Infants Not Admitted to Neonatal Intensive Care Unit. *Neonatology.* 2018;114(4):348-354. doi:[10.1159/000491817](https://doi.org/10.1159/000491817).
74. Kemper AR, Newman TB, Slaughter JL, et al. Clinical Practice Guideline Revision: Management of Hyperbilirubinemia in the Newborn Infant 35 or More Weeks of

Gestation. *Pediatrics*. 2022;150(3):e2022058859.
doi:[10.1542/peds.2022-058859](https://doi.org/10.1542/peds.2022-058859).

75. Sen S, Andrews C, Anderson E, Turner D, Monthé-Drèze C, Wachman EM. Type of feeding provided with dextrose gel impacts hypoglycemia outcomes: comparing donor milk, formula, and breastfeeding. *J Perinatol*. 2020;40(11):1705-1711.
doi:[10.1038/s41372-020-00776-y](https://doi.org/10.1038/s41372-020-00776-y).
76. Flaherman VJ, Aby J, Burgos AE, Lee KA, Cabana MD, Newman TB. Effect of Early Limited Formula on Duration and Exclusivity of Breastfeeding in At-Risk Infants: An RCT. *PEDIATRICS*. 2013;131(6):1059-1065.
doi:[10.1542/peds.2012-2809](https://doi.org/10.1542/peds.2012-2809)
77. Straňák Z, Feyereislova S, Černá M, Kollárová J, Feyereisl J. Limited Amount of Formula May Facilitate Breastfeeding: Randomized, Controlled Trial to Compare Standard Clinical Practice versus Limited Supplemental Feeding. *PLoS One*. 2016;11(2):1-12. doi:[10.1371/journal.pone.0150053](https://doi.org/10.1371/journal.pone.0150053).
78. Flaherman VJ, Narayan NR, Hartigan-O'Connor D, Cabana MD, McCulloch CE, Paul IA. The Effect of Early Limited Formula on Breastfeeding, Readmission, and Intestinal Microbiota: A Randomized Clinical Trial. *J Pediatr*. 2018;196:84-90.e1.
doi:[10.1016/j.jpeds.2017.12.073](https://doi.org/10.1016/j.jpeds.2017.12.073).
79. Kair LR, Flaherman VJ, Colaizy TT. Effect of Donor Milk Supplementation on Breastfeeding Outcomes in Term Newborns: A Randomized Controlled Trial. *Clin Pediatr*. 2019;58(5):534-540. doi:[10.1177/0009922819826105](https://doi.org/10.1177/0009922819826105).
80. Flaherman VJ, Cabana MD, McCulloch CE, Paul IM. Effect of Early Limited Formula on Breastfeeding Duration in the First Year of Life: A Randomized Clinical Trial. *JAMA Pediatr*. 2019;173(8):729-735.
doi:[10.1001/jamapediatrics.2019.1424](https://doi.org/10.1001/jamapediatrics.2019.1424).
81. Hoa DP, Thanh HT, Höjer B, Persson LA. Young child feeding in a rural area in the Red River delta, Vietnam. *Acta Paediatr*. 1995;84(9):1045-1049.
doi:[10.1111/j.1651-2227.1995.tb13822.x](https://doi.org/10.1111/j.1651-2227.1995.tb13822.x).
82. Srivastava SP, Sharma VK, Kumar V. Breast feeding pattern in neonates. *Indian Pediatr*. 1994;31(9):1079-1082. [PMID:7883365](https://pubmed.ncbi.nlm.nih.gov/7883365/).
83. Hasan J, Khan Z, Sinha SN. Socio-cultural factors influencing nutritional status of infants—a longitudinal study. *Indian J Matern Child Health*. 1991;2(3):84-86.
[PMID:12346054](https://pubmed.ncbi.nlm.nih.gov/12346054/).
84. Semega-Janneh IJ, Bøhler E, Holm H, Matheson I, Holmboe-Ottesen G. Promoting breastfeeding in rural Gambia: combining traditional and modern knowledge. *Health Policy Plan*. 2001;16(2):199-205. doi:[10.1093/heapol/16.2.199](https://doi.org/10.1093/heapol/16.2.199).

85. Davies-Adetugbo AA. Sociocultural factors and the promotion of exclusive breastfeeding in rural Yoruba communities of Osun State, Nigeria. *Soc Sci Med*. 1997;45(1):113-125. doi:[10.1016/s0277-9536\(96\)00320-6](https://doi.org/10.1016/s0277-9536(96)00320-6).
86. Delport SD, Bergh AM, Hay IT. Breast-feeding practices in a private maternity hospital. *S Afr Med J*. 1988;74(8):396-399. [PMID:3187818](https://pubmed.ncbi.nlm.nih.gov/3187818/).
87. Islam MA, Nielsen CC. Maternal and child health services: evaluating mothers' perceptions and participation. *Public Health*. 1993;107(4):243-249. doi:[10.1016/s0033-3506\(05\)80503-7](https://doi.org/10.1016/s0033-3506(05)80503-7).
88. Ashraf RN, Jalil F, Khan SR, et al. Early child health in Lahore, Pakistan: V. Feeding patterns. *Acta Paediatr Suppl*. 1993;82(Suppl 390):47-61. doi:[10.1111/j.1651-2227.1993.tb12906.x](https://doi.org/10.1111/j.1651-2227.1993.tb12906.x).
89. Sloand E, Budhathoki C, Junn J, Vo D, Lowe V, Pennington A. Breastfeeding among Latino families in an urban pediatric office setting. *Nurs Res Pract*. 2016;2016:9278401. doi:[10.1155/2016/9278401](https://doi.org/10.1155/2016/9278401).
90. Li R, Perrine CG, Anstey EH, Chen J, MacGowan CA, Elam-Evans LD. Breastfeeding trends by race/ethnicity among US children born from 2009 to 2015. *JAMA Pediatr*. 2019;173(12):e193319. doi:[10.1001/jamapediatrics.2019.3319](https://doi.org/10.1001/jamapediatrics.2019.3319).
91. Bhutani VK, Johnson LH. Newborn jaundice and kernicterus—health and societal perspectives. *Indian J Pediatr*. 2003;70(5):407-416. doi:[10.1007/BF02723615](https://doi.org/10.1007/BF02723615).